



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 684 240 A1**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: **95105495.6**

(22) Date of filing: **12.04.95**

(51) Int. Cl.<sup>6</sup>: **C07D 401/06, C07D 403/06,  
C07D 413/06, C07D 417/06,  
C07D 487/04, A61K 31/445,  
C07C 259/06**

(30) Priority: **25.04.94 GB 9408183  
30.01.95 GB 9501737**

(43) Date of publication of application:  
**29.11.95 Bulletin 95/48**

(64) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC  
NL PT SE**

(71) Applicant: **F. HOFFMANN-LA ROCHE AG  
Postfach 3255  
CH-4002 Basel (CH)**

(72) Inventor: **Broadhurst, Michael John  
Ivy Cottage,  
High Street**

**Barley,  
Royston, Hertfordshire (GB)  
Inventor: Brown, Paul Anthony**


**13 Highbury Road  
Hitchin,  
Hertfordshire (GB)**

**Inventor: Johnson, William Henry  
87 West Hill  
Hitchin,  
Hertfordshire (GB)**

(74) Representative: **Mezger, Wolfgang, Dr. et al  
Grenzacherstrasse 124  
Postfach 3255  
CH-4002 Basel (CH)**

(54) **Hydroxamic acid derivatives with tricyclic substitution.**

(57) The invention provides hydroxamic acid derivatives of the general formula



wherein R<sup>1</sup> represents cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl; R<sup>2</sup> represents a saturated 5- to 8-membered monocyclic or bridged N-heterocyclic ring which is attached via the N atom and which, when it is monocyclic, optionally contains NR<sup>4</sup>, O, S, SO or SO<sub>2</sub> as a ring member and/or is optionally substituted on one or more C atoms by hydroxy, lower alkyl, lower alkoxy, oxo, ketalized oxo, amino, mono(lower alkyl)amino, di-(lower alkyl)amino, carboxy, lower alkoxy carbonyl, hydroxy-methyl, lower alkoxy methyl, carbamoyl, mono(lower

**EP 0 684 240 A1**

alkyl)-carbamoyl, di(lower alkyl)carbamoyl or hydroxyimino; R<sup>3</sup> represents a 5- or 6-membered N-heterocyclic ring which (a) is attached via the N atom, (b) optionally contains N, O and/or S, SO or SO<sub>2</sub> as an additional ring member, (c) is substituted by oxo on one or both C atoms adjacent to the linking N atom and (d) is optionally benz-fused or optionally substituted on one or more other C atoms by lower alkyl or oxo and/or on any additional N atom(s) by lower alkyl or aryl; R<sup>4</sup> represents hydrogen, lower alkyl, aryl, aralkyl or a protecting group; m stands for 1 or 2 and n stands for 1-4, and pharmaceutically acceptable salts thereof, which are collagenase inhibitors useful in the control or prevention of degenerative joint diseases such as rheumatoid arthritis and osteoarthritis or in the treatment of invasive tumours, atherosclerosis or multiple sclerosis. They can be manufactured either by hydroxamidating a corresponding novel carboxylic acid or by deprotecting a corresponding novel benzyloxycarbamoyl compound.


The present invention is concerned with hydroxamic acid derivatives.

The hydroxamic acid derivatives provided by the present invention are compounds of the general formula

5

10

15



wherein

- 20     R¹    represents cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl;  
        R²    represents a saturated 5- to 8-membered monocyclic or bridged N-heterocyclic ring which is  
               attached via the N atom and which, when it is monocyclic, optionally contains NR⁴, O, S, SO or  
               SO₂ as a ring member and/or is optionally substituted on one or more C atoms by hydroxy, lower  
 25     alkyl, lower alkoxy, oxo, ketalized oxo, amino, mono(lower alkyl)amino, di(lower alkyl) amino,  
        carboxy, lower alkoxycarbonyl, hydroxymethyl, lower alkoxymethyl, carbamoyl, mono(lower alkyl)-  
        carbamoyl, di(lower alkyl)carbamoyl or hydroxyimino;  
        R³    represents a 5- or 6-membered N-heterocyclic ring which (a) is attached via the N atom, (b)  
               optionally contains N, O and/or S, SO or SO₂ as an additional ring member, (c) is substituted by  
               oxo on one or both C atoms adjacent to the linking N atom and (d) is optionally benz-fused or  
 30     optionally substituted on one or more other C atoms by lower alkyl or oxo and/or on any  
        additional N atom(s) by lower alkyl or aryl;  
        R⁴    represents hydrogen, lower alkyl, aryl, aralkyl or a protecting group;  
        m     stands for 1 or 2; and  
        n     stands for 1-4;

35   and pharmaceutically acceptable salts thereof.

The compounds of formula I possess valuable pharmacological properties. In particular, they are collagenase inhibitors and can be used in the control or prevention of degenerative joint diseases such as rheumatoid arthritis and osteoarthritis or in the treatment of invasive tumours, atherosclerosis or multiple sclerosis.

40   Objects of the present invention are the compounds of formula I and their pharmaceutically acceptable salts per se and for use as therapeutically active substances; a process for the manufacture of said compounds and salts; intermediates useful in said process; medicaments containing said compounds and salts and the manufacture of these medicaments; and the use of said compounds and salts in the control or prevention of illnesses or in the improvement of health, especially in the control or prevention of  
 45   degenerative joint diseases or in the treatment of invasive tumours or atherosclerosis, or for the manufacture of a medicament for the control or prevention of degenerative joint diseases or for the treatment of invasive tumours, atherosclerosis or multiple sclerosis.


As used in this Specification, the term "lower alkyl", alone or in combination, means a straight-chain or branched-chain alkyl group containing a maximum of six carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec.butyl, isobutyl, tert.butyl, n-pentyl, n-hexyl and the like. The term "lower alkoxy",  
 50   alone or in combination, means a straight-chain or branched-chain alkoxy group containing a maximum of six carbon atoms, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, tert.butoxy and the like. The term "aryl" means phenyl which is optionally substituted by, for example, lower alkyl, lower alkoxy and/or halogen, i.e. fluorine, chlorine, bromine or iodine, such as p-tolyl, p-methoxyphenyl, p-chlorophenyl and the  
 55   like. The term "aralkyl" means a lower alkyl group as hereinbefore defined in which one or more hydrogen atoms is/are replaced by an aryl group as hereinbefore defined, such as benzyl and the like. A ketalized oxo group can be, for example, ethylenedioxy.

A protecting group denoted by  $R^4$  can be any conventional protecting group, e.g. as known in peptide chemistry such as benzyloxycarbonyl, tert.butoxycarbonyl, acetyl and the like.

Examples of monocyclic N-heterocyclic rings denoted by  $R^2$  are 1-pyrrolidinyl, piperidino, 1-piperazinyl, 4-aryl-1-piperazinyl, hexahydro-1-pyridazinyl, morpholino, tetrahydro-1,4-thiazin-4-yl, tetrahydro-1,4-thiazin-4-yl 1-oxide, tetrahydro-1,4-thiazin-4-yl 1,1-dioxide, thiazolidin-3-yl, hexahydroazepino and octahydroazocino which can be substituted in the manner given earlier; for example 2-(methylcarbamoyl)-1-pyrrolidinyl, 2-(hydroxymethyl)-1-pyrrolidinyl, 4-hydroxypiperidino, 2-(methyl-carbamoyl)piperidino, 4-hydroxyiminopiperidino, 4-methoxypiperidino, 4-methyl-1-piperazinyl, 4-phenyl-1-piperazinyl, 1,4-dioxo-8-azaspiro-[4.5]decan-8-yl, hexahydro-3-(methylcarbamoyl)-2-pyridazinyl, hexahydro-1-(benzyloxycarbonyl)-2-pyridazinyl, 5,5-dimethyl-4-methylcarbamoyl-thiazolidin-3-yl and 5,5-dimethyl-4-propylcarbamoyl-thiazolidin-3-yl.

Examples of bridged N-heterocyclic rings denoted by  $R^2$  are 5-azabicyclo[2.1.1]hexane, 3-azabicyclo[3.1.1]heptane, 7-azabicyclo[2.2.1]heptane, 3-azabicyclo[3.2.1]octane, 2-azabicyclo[3.2.2]nonane and 3-azabicyclo[3.2.2]nonane.

Examples of N-heterocyclic rings denoted by  $R^3$  are rings of the formulae:



in which  $R^5$  and  $R^6$  each represent hydrogen or together represent an additional bond or the remainder of a fused benzene ring;

$R^7$  represents hydrogen, lower alkyl or aryl; and

- X represents -CO-, -CH<sub>2</sub>-, -CH(lower alkyl)-, -C(lower alkyl)<sub>2</sub>-, -NH-, -N(lower alkyl)- or -O-; or, when R<sup>7</sup> represents lower alkyl and X represents -N(lower alkyl)-, the lower alkyl groups can be joined to form a 5-, 6- or 7-membered ring;
- R<sup>8</sup> represents hydrogen, lower alkyl or aryl;
- 5 R<sup>9</sup> and R<sup>10</sup> each represent hydrogen or lower alkyl;
- Y represents -O-, -NH- or -N(lower alkyl)-; and
- Z represents S, SO or SO<sub>2</sub>;

Examples of such rings are 2-oxo-1-pyrrolidinyl, 2,5-dioxo-1-pyrrolidino, phthalimido, 1,2-dimethyl-3,5-dioxo-1,2,4-triazolidin-4-yl, 3-methyl-2,5-dioxo-1-imidazolidinyl, 3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl, 2-methyl-10 3,5-dioxo-1,2,4-oxadiazol-4-yl, 3-methyl-2,4,5-trioxo-1-imidazolidinyl, 2,5-dioxo-3-phenyl-1-imidazolidinyl, 2,6-dioxopiperidino, 5,5-dimethyl-2,4-dioxo-3-oxazolidinyl and hexahydro-1,3-dioxopyrazolo[1,2-a][1,2,4]-triazol-2-yl.

One group of preferred compounds of formula I comprises those in which R<sup>2</sup> represents 1-pyrrolidinyl, piperidino, 4-aryl-1-piperazino, morpholino, tetrahydro-1,4-thiazin-4-yl, tetrahydro-1,4-thiazin-4-yl 1,1-dioxide, 15 thiazolidin-3-yl, hexahydroazepino or octahydroazocino optionally substituted on one or more C atoms by hydroxy, lower alkyl, lower alkoxy, ketalized oxo or mono(lower alkyl)-carbamoyl, especially piperidino which is optionally substituted by hydroxy, particularly 4-hydroxypiperidino, or 3-azabicyclo[3.2.2]nonane. Also preferred are compounds of formula I in which R<sup>3</sup> represents a group of formula (b), (c), especially one in which R<sup>7</sup> represents lower alkyl and X represents -C(lower alkyl)<sub>2</sub>-, particularly 3,4,4-trimethyl-2,5-dioxo-20 1-imidazolidinyl, or (h). Preferably, m and n both stand for 1.

The most preferred compounds of formula I are:

- 1-[3-Cyclopropyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]piperidine,
- 1-[3-cyclopropyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]-4-piperidinol,
- 25 3-[3-cyclopropyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-3-azabicyclo[3.2.2]nonane,
- 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]piperidine,
- 30 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-4-piperidinol,
- 3-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-3-azabicyclo[3.2.2]nonane,
- 1-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-4-piperidinol,
- 35 3-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-3-azabicyclo[3.2.2]nonane and
- 1-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]piperidine.
- 40 Other preferred compounds of formula I hereinbefore are:
- 1-[3-cyclohexyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]piperidine,
- 4-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]tetrahydro-1,4-thiazine,
- 45 4-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]tetrahydro-1,4-thiazine S,S-dioxide,
- 4-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]tetrahydro-1,4-thiazine,
- 4-[3-cyclohexyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]tetrahydro-1,4-thiazine,
- 50 3-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-5,5-dimethyl-N-propyl-4(R)-thiazolidinecarboxamide,
- 4-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]morpholine,
- 55 3-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-N,5,5-trimethyl-4(R)-thiazolidinecarboxamide,
- 4-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-4-phenylpiperazine,


4-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]morpholine,  
 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]pyrrolidine,  
 5 8-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-1,4-dioxo-8-azaspiro[4,5]decane,  
 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-4-methoxypiperidine,  
 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-  
 10 propionyl]octahydroazocine,  
 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(5,5-dimethyl-2,4-dioxo-3-oxazolidinyl)ethyl]-propionyl]piperidine,  
 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]hexahydroazepine,  
 15 1-[3-cyclobutyl-2(R)-[2-(hexahydro-1,3-dioxypyrazolo[1,2-a][1,2,4]triazol-2-yl)-1(R or S)-(hydroxycarbamoyl)ethyl]propionyl]piperidine and  
 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-phthalimidoethyl]propionyl]piperidine.

The compounds of formula I form pharmaceutically acceptable salts with bases such as alkali metal hydroxides (e.g. sodium hydroxide and potassium hydroxide), alkaline earth metal hydroxides (e.g. calcium  
 20 hydroxide and magnesium hydroxide), ammonium hydroxide and the like. The compounds of formula I which are basic form pharmaceutically acceptable salts with acids. As such salts there come into consideration not only salts with inorganic acids such as hydrohalic acids (e.g. hydrochloric acid and hydrobromic acid), sulphuric acid, nitric acid, phosphoric acid etc, but also salts with organic acids such as acetic acid, tartaric acid, succinic acid, fumaric acid, maleic acid, malic acid, salicylic acid, citric acid,  
 25 methanesulphonic acid, p-toluenesulphonic acid etc.

The compounds of formula I contain at least two asymmetric carbon atoms and can accordingly exist as optically active enantiomers, as diastereoisomers or as racemates. The present invention is intended to embrace all of these forms.

According to the process provided by the present invention, the compounds of formula I and their  
 30 pharmaceutically acceptable salts are manufactured by

(a) reacting an acid of the general formula




wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, m and n have the significance given earlier, with a compound of the general  
 formula

50 H<sub>2</sub>N-OZ (III)

wherein Z represents hydrogen, tri(lower alkyl)silyl or diphenyl(lower alkyl)silyl,  
 and, where required, cleaving off any diphenyl(lower alkyl)silyl group present in the reaction product,  
 or

55 (b) catalytically hydrogenating a compound of the general formula



15 wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $m$  and  $n$  have the significance given earlier and Bz represents benzyl, and,

if desired, converting a compound of formula I obtained into a pharmaceutically acceptable salt.


The reaction of an acid of formula II with a compound of formula III in accordance with embodiment (a) of the process can be carried out in a known manner. For example, an acid of formula II can be reacted  
20 with a compound of formula III in an inert organic solvent such as dichloromethane, dimethylformamide or the like using 1-hydroxybenzotriazole in the presence of a condensation agent such as 1-ethyl-3-(3-dimethylamino-propyl)carbodiimide hydrochloride at about 0°C to about room temperature. Alternatively, an acid of formula II can be converted into the corresponding acid chloride (e.g. using oxalyl chloride) and the acid chloride can then be reacted with a compound of formula III. Preferred compounds of formula III are  
25 those in which Z represents tert.butyl dimethylsilyl or tert.butyl diphenylsilyl. When a compound of formula III in which Z represents tri(lower alkyl)silyl is used, this group is cleaved off during the reaction and working-up, and a compound of formula I is obtained directly. On the other hand, when a compound of formula III in which Z represents diphenyl(lower alkyl)silyl is used, this group remains in the reaction product and must subsequently be cleaved off in a known manner, for example by means of fluoride ions.

30 The catalytic hydrogenation of a compound of formula IV in accordance with embodiment (b) of the process can be carried out in a manner known per se; for example in an inert organic solvent using hydrogen in the presence of a noble metal catalyst. Suitable inert organic solvents are, for example, lower alkanols such as methanol, ethanol, etc. With respect to the catalyst, this can be, for example, a platinum, palladium or rhodium catalyst which can be supported on a suitable carrier material. Palladium-on-charcoal  
35 is the preferred catalyst. The temperature and pressure are not critical, although for convenience the catalytic hydrogenation is preferably carried out at room temperature and under atmospheric pressure.

Compounds of formula I can be converted into pharmaceutically acceptable salts by treatment with bases and basic compounds of formula I can be converted into pharmaceutically acceptable salts by treatment with acids. Such treatments can be carried out in a conventional manner.

40 The acids of formula II which are used as starting materials in embodiment (a) of the process are novel and form a further object of the present invention.

The acids of formula II can be prepared, for example, as illustrated in the following Reaction Scheme in which  $R^1$ ,  $R^2$ ,  $R^3$ ,  $m$  and  $n$  have the significance given earlier, Bz represents benzyl and tBu represents tert-butyl:

Reaction Scheme

Having regard to the foregoing Reaction Scheme, the individual steps thereof can be carried out according to methods known per se. Thus, in the first step, an amino acid of formula V, which can be obtained according to the procedure described by Chenault H.K, Dahmer J. and Whitesides G.M., J.Am.Chem.Soc. 1989, 111, 6354-6364, is converted by treatment with sodium nitrite in the presence of concentrated sulphuric acid into a hydroxy acid of formula VI which is subsequently reacted with benzyl bromide in the presence of an organic base, e.g. a trialkylamine such as triethylamine, into a corresponding



benzyl ester of formula VII. The latter is then activated, e.g. by reaction with trifluoromethanesulphonic anhydride, and treated with benzyl tert-butyl malonate in the presence of a strong base, e.g. an alkali metal hydride such as sodium hydride, to give a compound of formula VIII. Treatment of the latter with a strong base, e.g. an alkali metal hydride such as sodium hydride, and reaction with a compound of formula IX yields a dibenzyl tert-butyl butanetricarboxylate of formula X which is then debenzylated by catalytic hydrogenation, e.g. in the presence of a palladium catalyst such as palladium-on-charcoal, to give a tert-butyl dihydrogen butanetricarboxylate of formula XI. Decarboxylation of this compound, e.g. by heating in toluene with triethylamine, which may be carried out in situ, yields a tert-butyl hydrogen succinate of formula XII which is condensed with a cyclic amine of formula XIII, e.g. according to the acid chloride method or using 1-hydroxybenzotriazole in the presence of a condensation agent such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, to give a compound of formula XIV which is deprotected (e.g. by treatment with trifluoroacetic acid) to give an acid of formula II.

The compounds of formula IV which are used as starting materials in embodiment (b) of the process are novel and form a further object of the present invention.

The compounds of formula IV can be prepared, for example, by reacting an acid of formula II with O-benzylhydroxylamine. This reaction can be carried out in a known manner, for example in an inert organic solvent such as dichloromethane or dimethylformamide using 1-hydroxybenzotriazole in the presence of a condensation agent such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride.

The remaining compounds which are used as intermediates or reactants in the manufacture of the compounds of formula I are known compounds or analogues of known compounds which can be prepared in a similar manner to the known compounds.

As mentioned earlier, the compounds of formula I and their pharmaceutically acceptable salts are collagenase inhibitors. The in vitro collagenase inhibiting activity of the present compounds and salts can be demonstrated using collagenase obtained from a culture of human synovial fibroblasts according to the method of Dayer J-M et al., Proc. Natl. Acad. Sci. USA (1976), 73 945, following activation of the pro-collagenase in the conditioned medium by treatment with trypsin. Collagenase activity was measured using <sup>14</sup>C-acetylated collagen type I from rat tail tendons as the substrate and employing the microtitre plate assay method of Johnson-Wint, B, Anal. Biochem. (1980), 104, 175. The IC<sub>50</sub> is that concentration of a compound or salt of the present invention in the enzyme digestion which reduces substrate cleavage and solubilization to 50% of that achieved by the enzyme alone.

The results obtained in the foregoing test with representative compounds and salts of this invention are compiled in Table I hereinafter:

Table I

Product of Example No.	IC <sub>50</sub> (nM)
2	18.0
4	7.0
5	2.5
7	6.5
9	8.5
16	4.1
17	2.35
23	34.0

The compounds of formula I and their pharmaceutically acceptable salts can be used as medicaments, for example in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered orally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions. However, they can also be administered rectally, e.g. in the form of suppositories, or parenterally, e.g. in the form of injection solutions.

For the manufacture of pharmaceutical preparations the compounds of formula I and their pharmaceutically acceptable salts can be formulated with therapeutically inert, inorganic or organic carriers. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts can be used, for example, as such carriers for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carriers for soft gelatine capsules are, for example, vegetable oils, waxes, fats, semi-solid and liquid polyols and the like. Depending on the nature of the active ingredient no carriers are, however, generally required in the case of soft gelatine capsules.

Suitable carriers for the manufacture of solutions and syrups are, for example, water, polyols, saccharose, invert sugar, glucose and the like. Suitable carriers for the manufacture of injection solutions are, for example, water, alcohols, polyols, glycerine, vegetable oils and the like. Natural and hardened oils, waxes, fats, semi-liquid polyols and the like are suitable carriers for the manufacture of suppositories.

- 5 The pharmaceutical preparations can also contain preservatives, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for adjustment of the osmotic pressure buffers coating agents or antioxidants.

Medicaments containing a compound of formula I or a pharmaceutically acceptable salt thereof and a therapeutically acceptable carrier as well as a process for the manufacture of such medicaments are also  
10 objects of the present invention. This process comprises mixing a compound of formula I or a pharmaceutically acceptable salt thereof with a therapeutically inert carrier material and bringing the mixture into a galenical administration form.

As mentioned earlier, the compounds of formula I and their pharmaceutically acceptable salts can be used in the control or prevention of illnesses, especially in the control or prevention of degenerative joint  
15 diseases or in the treatment of invasive tumours, atherosclerosis or multiple sclerosis. The dosage can vary within wide limits and will, of course, be adjusted to the individual requirements in each particular case. In general, in the case of administration to adults, a daily dosage of from about 5 mg to about 30 mg, preferably from about 10 mg to about 15 mg, should be appropriate, although the upper limit may be exceeded when this is found to be expedient. The daily dosage can be administered as a single dosage or  
20 in divided dosages.

The following Examples illustrate the present invention in more detail. In these Examples all temperatures are given in degrees Celsius.

#### Example 1

25 A solution of 0.575 g of 1-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopropylpropionyl]piperidine (diastereoisomer 1) in 10 ml of ethanol was hydrogenated in the presence of 0.4 g of 5% palladium-on-charcoal catalyst for 6 hours. The catalyst was removed by filtration and the solution was evaporated. The residue was purified by flash chromatography on silica  
30 gel using dichloromethane/methanol (96:4) for the elution to give 0.37 g of 1-[3-cyclopropyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]piperidine (diastereoisomer 1) in the form of a white foam. nmr (MeOD): 3.78-3.64 (m, 3H); 3.62 (dd, 1H, J = 15,8); 3.49-3.41 (m, 1H); 3.39 (dd, 1H, J = 15,5); 3.33-3.27 (m, 1H); 2.95-2.87 (m, 1H); 2.83 (s, 3H); 1.74-1.46 (m, 7H); 1.33 (s, 3H); 1.31 (s, 3H); 1.20-1.13 (m, 1H); 0.61-0.50 (m, 1H); 0.44-0.33 (m, 2H); 0.06- -0.05 (m, 2H);  
35 MS: 409 (M + H)<sup>+</sup>.

The starting material was prepared as follows:

(i) A solution of 4.9 g of 2(R)-amino-3-cyclopropylpropionic acid (prepared in a manner analogous to that  
40 described by Chenault H.K., Dahmer J. and Whitesides G.M. in J. Am. Chem. Soc. 1989, 111, 6354-6364) in 50 ml of water containing 4.05 ml of concentrated sulphuric acid was warmed to 45°. A solution of 10.5 g of sodium nitrite in 20 ml of water was added dropwise over 30 minutes. The solution was stirred at 45° for 4 hours and then cooled to room temperature. The solution was extracted with three 50 ml portions of ethyl acetate. The combined extracts were washed with water and dried over anhydrous  
45 magnesium sulphate. The solvent was evaporated to leave 3.95 g of a yellow oil containing 3-cyclopropyl-2(R)-hydroxypropionic acid which was used in the next step without further purification.

Rf [dichloromethane/methanol (9:1)] = 0.65.

(ii) A solution of 3.95 g of the product from (i) in 50 ml of ethyl acetate was treated with 5.32 ml of triethylamine and 3.8 ml of benzyl bromide. The mixture was stirred and heated under reflux for 3 hours,  
50 then allowed to cool to room temperature overnight. The suspension was washed with 2M hydrochloric acid, water and saturated sodium chloride solution. After drying over anhydrous magnesium sulphate the solvent was evaporated. The residue was purified by flash chromatography on silica gel using hexane/ethyl acetate (2:1) for the elution to give 3.36 g of benzyl 3-cyclopropyl-2(R)-hydroxypropionate in the form of a yellow oil.

55 nmr (CDCl<sub>3</sub>): 7.39-7.28 (m, 5H); 5.19 (d, 1H, J = 14); 5.15 (d, 1H, J = 14); 4.31-4.24 (m, 1H); 2.81 (br. d, 1H); 1.69-1.54 (m, 2H); 0.87-0.74 (m, 1H); 0.45-0.34 (m, 2H); 0.08- -0.07 (m, 2H).

(iii) A solution of 3.36 g of the product from (ii) and 1.49 ml of pyridine in 10 ml of dichloromethane was added dropwise to a solution of 3.07 ml of trifluoromethanesulphonic anhydride in 15 ml of dich-

loromethane at 0° over 30 minutes with stirring. The mixture was stirred at 0° for 2 hours and then washed with water and saturated sodium chloride solution. After drying over anhydrous magnesium sulphate the solvent was evaporated to give 5.37 g of benzyl 3-cyclopropyl-2(R)-trifluoromethylsulphonyloxypropionate in the form of an orange oil which was used in the next step without further purification.

$R_f$ [hexane/ethyl acetate (4:1)] = 0.5.

(iv) A solution of 3.8 g of benzyl tert-butyl malonate in 50 ml of 1,2-dimethoxyethane was treated with 0.504 g of an 80% dispersion of sodium hydride in mineral oil. The mixture was stirred at room temperature for 30 minutes and then cooled to 0°. A solution of 5.37 g of the product from (iii) in 20 ml of dichloromethane was added dropwise at 0°. The mixture was stirred at 0° for 2 hours and then left to warm to room temperature overnight. The solvent was evaporated and the residue was dissolved in ethyl acetate. The solution was washed with water and saturated sodium chloride solution. After drying over anhydrous magnesium sulphate the solvent was evaporated to give 6.54 g of 2,3-dibenzyl 3-tert-butyl 1-cyclopropyl-2(R),3(R,S),3-propanetricarboxylate as a 1:1 mixture of diastereoisomers in the form of an orange oil.

nmr (CDCl<sub>3</sub>): 7.46-7.36 (m, 20H); 5.19-5.07 (m, 8H); 3.89 (d, 1H, J = 10); 3.85 (d, 1H, J = 10) 3.37-3.26 (m, 2H); 1.68-1.52 (m, 2H); 1.52-1.38 (m, 2H); 1.41 (s, 9H); 1.39 (s, 9H); 0.79-0.63 (m, 2H); 0.49-0.38 (m, 4H); 0.12-0.07 (m, 4H).

(v) A solution of 6.4 g of the product from (iv) in 30 ml of 1,2-dimethoxyethane was treated with 0.446 g of an 80% dispersion of sodium hydride in mineral oil. The mixture was stirred at room temperature for 30 minutes. A solution of 3.84 g of 1-(bromomethyl)-3,4,4-trimethyl-2,5-imidazolidinedione in 20 ml of 1,2-dimethoxyethane was added dropwise over 15 minutes. The mixture was stirred at room temperature for 36 hours, the solvent was evaporated and the residue was dissolved in ethyl acetate and washed with water and saturated sodium chloride solution. After drying over anhydrous magnesium sulphate the solvent was evaporated. The residue was purified by flash chromatography on silica gel using hexane/ethyl acetate (7:3) and subsequently hexane/ethyl acetate (6:4) for the elution to give 6.4 g of 2,3-dibenzyl 3-tert-butyl 1-cyclopropyl-4-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-2(R),3(R,S),3-butanetricarboxylate as a 1:1 mixture of diastereoisomers in the form of a clear oil.

nmr (CDCl<sub>3</sub>): 7.47-7.28 (m, 20H); 5.31-5.03 (m, 8H); 4.32-4.18 (m, 4H); 3.19-3.15 (m, 1H); 3.16-3.12 (m, 1H); 2.86 (s, 6H); 2.00-1.90 (m, 1H); 1.89-1.79 (m, 1H); 1.64-1.49 (m, 1H); 1.48-1.38 (m, 1H); 1.37 (s, 12H); 1.36 (s, 9H); 1.32 (s, 9H); 0.9-0.8 (m, 2H); 0.41-0.3 (m, 4H); 0.15-0.05 (m, 2H); 0.04-0.04 (m, 2H).

(vi) A solution of 3.0 g of the product from (v) in 30 ml of 2-propanol was hydrogenated in the presence of 0.3 g of 5% palladium on charcoal catalyst for 2 hours. The catalyst was removed by filtration and the solution was evaporated. The residue was re-evaporated from 20 ml toluene and then dissolved in 50 ml of toluene. The solution was treated with 0.693 ml of triethylamine and the mixture was heated under reflux for 2 hours. The solution was cooled to room temperature and washed with 2M hydrochloric acid, water and saturated sodium chloride solution. After drying over anhydrous magnesium sulphate the solvent was evaporated to give 1.85 g of 4-tert-butyl hydrogen 2(R)-(cyclopropylmethyl)-3(R or S)-[(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)methyl]succinate as an approximately 6:1 mixture of diastereoisomers in the form of a yellow oil.

MS: 383 (M + H)<sup>+</sup>;

$R_f$  [dichloromethane/methanol (9:1)] = 0.41.

(vii) A solution of 1.0 g of the product from (vi) in 10 ml of dichloromethane was cooled to 0° and treated in succession with 0.665 ml of N-ethylmorpholine, 0.481 g of 1-hydroxybenzotriazole and 0.602 g of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride. The mixture was stirred at 0° for 30 minutes and then treated with 0.517 ml of piperidine. The solution was left to warm to room temperature and was stirred overnight. The solution was washed with 5% aqueous sodium hydrogen carbonate solution, 2M hydrochloric acid and saturated sodium chloride solution. After drying over anhydrous magnesium sulphate the solvent was evaporated to give 1.01 g of 1-[2(R)-[1(R or S)-(tert-butoxycarbonyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopropylpropionyl]piperidine as an approximately 6:1 mixture of diastereoisomers in the form of a yellow gum.

MS: 450 (M + H)<sup>+</sup>;

$R_f$  [dichloromethane/methanol (95:5)] = 0.51.

(viii) A solution of 1.0 g of the product from (vii) in 2 ml of trifluoroacetic acid was stirred at room temperature for 2.5 hours. The solvent was evaporated and the residue was re-evaporated from toluene. The residue was dissolved in diethyl ether and the solution was extracted with two portions of 5% aqueous sodium hydrogen carbonate solution. The combined extracts were acidified to pH 2 with concentrated hydrochloric acid and the product was extracted with two portions of dichloromethane. The

combined organic extracts were washed with water and saturated sodium chloride solution and dried over anhydrous magnesium sulphate. The solvent was evaporated to give 0.634 g of a white foam containing 1-[2(R)-[1(R or S)carboxy-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopropylpropionyl]piperidine as a 6:1 mixture of diastereoisomers which was used in the next step without further purification.

Rf [dichloromethane/methanol (9:1)] = 0.31.

(ix) A solution of 0.634 g of the product from (viii) in 10 ml of dichloromethane was cooled at 0°. The solution was treated in succession with 0.41 ml of N-ethylmorpholine, 0.296 g of 1-hydroxybenzotriazole and 0.371 g of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride. The mixture was stirred at 0° for 30 minutes. A solution of 0.238 g of O-benzylhydroxylamine in 2 ml of dichloromethane was added. The mixture was left to warm to room temperature and was stirred overnight. The solution was washed with two portions of 5% aqueous sodium hydrogen carbonate solution and subsequently with 2M hydrochloric acid, water and saturated sodium chloride solution. After drying over anhydrous magnesium sulphate, the solvent was removed by evaporation. The residue was purified by flash chromatography on silica gel using dichloromethane/methanol (98:2) for the elution to give 0.592 g of 1-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopropylpropionyl]-piperidine (diastereoisomer 1) as a white foam.

nmr (MeOD): 7.45-7.31 (m, 5H); 4.87 (d, 1H, J = 13); 4.79 (d, 1H, J = 13); 3.78-3.65 (m, 3H); 3.63 (dd, 1H, J = 15, 8); 3.53-3.45 (m, 1H); 3.44 (dd, 1H, J = 15, 5); 3.34-3.27 (m, 1H); 2.87 (s, 3H); 2.84-2.78 (m, 1H); 1.78-1.49 (m, 7H); 1.49-1.40 (m, 1H); 1.36 (s, 3H); 1.32 (s, 3H); 1.12-1.04 (m, 1H); 0.61-0.50 (m, 1H); 0.48-0.37 (m, 2H); 0.07-0.06 (m, 2H).

MS: 499 (M + H)<sup>+</sup>.

#### Example 2

In a manner analogous to that described in the first paragraph of Example 1, from 0.391 g of 1-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopropylpropionyl]-4-piperidinol (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.33 g of 1-[3-cyclopropyl-2-(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-4-piperidinol (diastereoisomer 1) in the form of a white foam.

nmr (MeOD): 4.22-4.02 (m, 2H); 3.90-3.81 (m, 1H); 3.69-3.56 (m, 1H); 3.49-3.38 (m, 2H); 3.37-3.18 (m, 2H); 3.11-3.01 (m, 1H); 2.97-2.86 (m, 1H); 2.83 (d, 3H, J = 5); 2.01-1.78 (m, 2H); 1.68-1.36 (m, 3H); 1.33 (s, 3H); 1.31 (d, 3H, J = 5); 1.24-1.13 (m, 1H); 0.62-0.50 (m, 1H); 0.49-0.33 (m, 2H); 0.09-0.05 (m, 2H);

MS: 425 (M + H)<sup>+</sup>

#### Example 3

In a manner analogous to that described in the first paragraph of Example 1, from 0.822 g of 3-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopropyl]-3-azabicyclo[3.2.2]nonane (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.496 g of 3-[3-cyclopropyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-3-azabicyclo[3.2.2]nonane (diastereoisomer 1) in the form of a white foam.

nmr (MeOD): 4.0-3.1 (m, 5H); 3.48-3.31 (m, 2H); 2.96-2.86 (m, 1H); 2.82 (s, 3H); 2.14-2.03 (m, 2H); 1.80-1.68 (m, 4H); 1.68-1.53 (m, 5H); 1.32 (s, 3H); 1.31 (s, 3H); 1.21-1.12 (m, 1H); 0.64-0.52 (m, 1H); 0.45-0.33 (m, 2H); 0.08-0.05 (m, 2H);

MS: 449 (M + H)<sup>+</sup>.

#### Example 4

In a manner analogous to that described in the first paragraph of Example 1, from 0.6 g of 1-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-piperidine (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.5 g of 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]piperidine (diastereoisomer 1) in the form of a white foam.

nmr (MeOD): 3.67 (dd, 1H, J = 15, 10); 3.64-3.46 (m, 4H); 3.34 (dd, 1H, J = 15, 8); 3.12 (td, 1H, J = 13, 3); 2.92-2.84 (m, 1H); 2.82 (s, 3H); 2.22-2.09 (m, 1H); 2.07-1.93 (m, 2H); 1.90-1.42 (m, 12H); 1.33 (s, 3H); 1.32 (s, 3H);

MS: 423 (M + H)<sup>+</sup>.

#### Example 5

- 5 In a manner analogous to that described in the first paragraph of Example 1, from 0.4 g of 1-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-4-piperidinol (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.294 g of 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-4-piperidinol in the form of a white foam.
- 10 nmr (MeOD): 4.15-4.05 (m, 1H); 4.04-3.90 (m, 1H); 3.90-3.80 (m, 1H); 3.72-3.57 (m, 1H); 3.45-3.30 (m, 2H); 3.18-3.06 (m, 2H); 2.94-2.85 (m, 1H); 2.84 (d, 3H, J = 5); 2.21-1.36 (m, 13H); 1.33 (d, 3H, J = 3); 1.31 (d, 3H, J = 6);
- MS: 439 (M + H)<sup>+</sup>.

#### Example 6

- 15 In a manner analogous to that described in the first paragraph of Example 1, from 0.642 g of 3-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutyl]-3-azabicyclo[3.2.2]nonane (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.348 g of 3-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-3-azabicyclo[3.2.2]nonane (diastereoisomer 1) in the form of a white foam.
- 20 nmr (MeOD): 3.92-3.83 (m, 2H); 3.76 (dd, 1H, J = 15, 13); 3.67-3.57 (m, 2H); 3.34 (dd, 1H, J = 15, 5); 3.28-3.21 (m, 1H); 2.96-2.87 (m, 1H); 2.83 (s, 3H); 2.23-2.13 (m, 1H); 2.12-1.92 (m, 4H); 1.91-1.48 (m, 14H); 1.35 (s, 3H); 1.34 (s, 3H).
- 25 MS: 463 (M + H)<sup>+</sup>.

#### Example 7

- 30 In a manner analogous to that described in the first paragraph of Example 1, from 0.5 g of 1-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopentylpropionyl]-4-piperidinol (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.4 g of 1-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-4-piperidinol (diastereoisomer 1) in the form of a white foam.
- 35 nmr (MeOD): 4.20-4.02 (m, 2H); 3.91-3.83 (m, 1H); 3.76-3.64 (m, 1H); 3.48-3.32 (m, 2H); 3.26-3.08 (m, 3H); 2.05-1.42 (m, 12H); 1.38-1.25 (m, 7H); 1.18-1.01 (m, 3H);
- MS: 453 (M + H)<sup>+</sup>.

#### Example 8

- 40 In a manner analogous to that described in the first paragraph of Example 1, from 0.57 g of 3-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopentyl]-3-azabicyclo[3.2.2]nonane (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.48 g of 3-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-3-azabicyclo[3.2.2]nonane (diastereoisomer 1) in the form of a white foam.
- 45 nmr (MeOD): 3.88-3.67 (m, 5H); 3.39-3.31 (m, 2H); 2.92-2.85 (m, 4H); 2.15-2.06 (m, 2H); 1.83 - 1.45 (m, 16H); 1.36-1.28 (m, 7H); 1.16-1.02 (m, 2H).
- MS: 477 (M + H)<sup>+</sup>.

#### Example 9

- 50 A solution of 0.421 g of an approximately 6:1 mixture of diastereoisomer 1 and diastereoisomer 2 of 1-[2(R)-[1(R or S)-carboxy-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopentylpropionyl]-piperidine, prepared in a manner analogous to that described in Example 1(i)-(viii), in 10 ml of dichloromethane was cooled to 0°. The solution was treated with 0.211 g of 1-hydroxybenzotriazole, 0.24 g of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride and 0.22 ml of N-methylmorpholine. The mixture was stirred at 0° for 15 minutes. A solution of 0.295 g of O-(tert-butyl dimethylsilyl)hydroxylamine and

## EP 0 684 240 A1

0.22 ml of N-methylmorpholine in 5 ml of dichloromethane was added. The mixture was left to warm to room temperature and was stirred overnight. The solution was washed with two portions of 5% aqueous sodium hydrogen carbonate solution and subsequently with 2M hydrochloric acid and saturated sodium chloride solution. After drying over anhydrous magnesium sulphate, the solvent was evaporated. The residue was purified by flash chromatography on silica gel using dichloromethane/ methanol (96:4) for the elution to give 0.123 g of 1-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]piperidine (diastereoisomer 1) in the form of a white foam.

nmr (MeOD): 3.74-3.66 (m, 3H); 3.53-3.45 (m, 2H); 3.34 (dd, J = 14,7,1H); 3.23 (dt, J = 4, 14, 1H); 2.90-2.84 (m, 4H); 1.80-1.45 (m, 14H); 1.38-1.23 (m, 7H); 1.15-1.01 (m, 2H);

MS: 437 (M + H)<sup>+</sup>.

### Example 10

In a manner analogous to that described in the first paragraph of Example 1, starting from 0.328 g of 1-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclohexylpropionyl]piperidine (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.269 g of 1-[3-cyclohexyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]piperidine (diastereoisomer 1) in the form of a white foam.

nmr (MeOD): 3.87-3.77 (m, 2H); 3.7 (dd, J = 14,9,1H); 3.64-3.56 (m, 2H); 3.38-3.28 (m, 2H); 2.9-2.83 (m, 4H); 1.84-1.45 (m, 12H); 1.35(s, 3H); 1.33 (s, 3H); 1.25-1.05 (m, 5H); 0.98-0.78 (m, 2H).

MS: 451 (M + H)<sup>+</sup>.

### Example 11

In a manner analogous to that described in Example 9, starting from 0.8 g of 1-[2(R)-[1(R or S)-carboxy-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopentylpropionyl]-tetrahydro-1,4-thiazine (diastereoisomer 1), prepared in a manner analogous to Example 1 (i)-(viii), there was obtained 0.3 g of 4-[3-cyclopentyl-2(R)-[1-(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-tetrahydro-1,4-thiazine (diastereoisomer 1) in the form of a white foam.

nmr (MeOD): 4.02-3.96 (m, 2H); 3.92-3.85 (m, 2H); 3.7 (dd, J = 13,9,1H); 3.37 (dd, J = 13,6,1H); 3.25-3.18 (m, 1H); 2.9-2.84 (m, 4H); 2.82-2.75 (m, 1H); 2.7-2.55 (m, 3H); 1.78-1.45 (m, 8H); 1.35 (s, 3H); 1.34 (s, 3H); 1.18-1.04 (m, 2H).

MS: 455 (M + H)<sup>+</sup>.

### Example 12

In a manner analogous to that described in Example 1, starting from 0.3 g of 4-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopentylpropionyl]-tetrahydro-1,4-thiazine S,S-dioxide (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.2 g of 4-[3-cyclopentyl-2(R)-[1-(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-tetrahydro-1,4-thiazine S,S dioxide (diastereoisomer 1) in the form of a white solid.

nmr (MeOD): 4.45-4.3 (m, 2H); 4.0-3.93 (m, 1H); 3.78-3.65 (m, 2H); 3.55-3.39 (m, 2H); 3.30-3.21 (m, 2H); 3.14-3.03 (m, 2H); 2.9-2.85 (m, 4H); 1.78-1.45 (m, 9H); 1.36 (s, 3H); 1.34 (s, 3H); 1.18-1.0 (m, 2H).

MS: 487 (M + H)<sup>+</sup>.

### Example 13

In a manner analogous to that described in Example 9, starting from 0.8 g of 1-[2(R)-[1(R or S)-carboxy-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-tetrahydro-1,4-thiazine (diastereoisomer 1), prepared in a manner analogous to Example 1 (i)-(viii), there was obtained 0.24 g of 4-[3-cyclobutyl-2(R)-[1-(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-tetrahydro-1,4-thiazine (diastereoisomer 1) in the form of a white solid.

nmr (MeOD): 3.98-3.75 (m, 4H); 3.64 (dd, J = 13,8,1H); 3.35 (dd, J = 15,6,1H); 3.07 (td, J = 10,4,1H); 2.9-2.83 (m, 1H); 2.82 (s, 3H); 2.78-2.72 (m, 1H); 2.66-2.52 (m, 3H); 2.18-2.08 (m, 1H); 2.05-1.93 (m, 2H); 1.85-1.45 (m, 6H); 1.13 (s, 3H); 1.11 (s, 3H).

MS: 441 (M + H)<sup>+</sup>.

Example 14

In a manner analogous to that described in Example 9, starting from 1.22 g of 1-[2(R)-[1-(R or S)-carboxy-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclohexylpropionyl]-tetrahydro-1,4-thiazine (diastereoisomer 1), prepared in a manner analogous to Example 1 (i)-(viii), there was obtained 0.45 g of 4-[3-cyclohexyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-tetrahydro-1,4-thiazine (diastereoisomer 1) in the form of a white solid.

nmr (MeOD): 4.12-4.03 (m, 2H); 3.95-3.88 (m, 1H); 3.75-3.65 (m, 2H); 3.38 (dd, J = 14,6,1H); 2.88-2.82 (m, 4H); 2.78-2.72 (m, 1H); 2.68-2.55 (m, 3H); 1.82-1.53 (m, 7H); 1.35 (s, 3H); 1.34 (s, 3H); 1.26-0.8 (m, 8H);

MS: 469 (M + H)<sup>+</sup>.

Example 15

In a manner analogous to that described in Example 9, from 1.164 g of a mixture of diastereoisomers of 3-[2(R)-[1(RS)-carboxy-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopentylpropionyl]-5,5-dimethyl-N-propyl-4(R)-thiazolidinecarboxamide, prepared in a manner analogous to that described in Example 1 (i)-(viii), there was obtained 0.329 g of 3-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-5,5-dimethyl-N-propyl-4(R)-thiazolidinecarboxamide (diastereoisomer 1) in the form of a white solid.

nmr (MeOD): 5.09-4.72 (m, 2H); 4.51 and 4.46 (both s, total 1H); 3.84 and 3.64 (both dd, J = 14,8,1H); 3.40-3.05 (m, 4H); 2.90-2.73 (m, 4H); 1.94-1.25 (m, 23H); 1.23-1.01 (m, 2H); 0.99-0.85 (m, 3H);

MS: 554 (M + H)<sup>+</sup>.

Example 16

In a manner analogous to that described in the first paragraph of Example 1, from 0.223 g of 4-[2(R)-[R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopentylpropionyl]-morpholine (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.112 g of 4-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]morpholine (diastereoisomer 1) in the form of a white solid.

nmr (MeOD): 3.83-3.56 (m, 9H); 3.41 (dd, J = 14,6, 1H); 3.19 (dt, J = 4,11,1H); 2.91-2.81 (m, 4H); 1.77-1.42 (m, 8H); 1.38-1.23 (m, 7H); 1.19-0.99 (m, 2H);

MS: 439 (M + H)<sup>+</sup>.

Example 17

In a manner analogous to that described in Example 9, from 1.289 g of a mixture of diastereoisomers of 3-[2(R)-[1(RS)-carboxy-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopentylpropionyl]-N,5,5-trimethyl-4(R)-thiazolidinecarboxamide (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(viii), there was obtained 0.629 g of 3-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-N,5,5-trimethyl-4(R)-thiazolidinecarboxamide (diastereoisomer 1) in the form of a white solid.

nmr (MeOD): 4.09-4.51 (m, 2H); 4.47 and 4.43 (both s, total 1H); 3.82 and 3.62 (both dd, J = 14,10, total 1H); 3.37 and 3.17 (both dd, J = 14,5, total 1H); 3.13-2.70 (m, 8H); 1.96-1.25 (m, 21H); 1.23-0.99 (m, 2H);

MS: 526 (M + H)<sup>+</sup>.

Example 18

In a manner analogous to that described in the first paragraph of Example 1, from 0.289 g of 1-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-4-phenylpiperazine (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.121 g of 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)methyl]-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-4-phenylpiperazine (diastereoisomer 1) in the form of a white solid.

nmr (MeOD): 7.25 (m, 2H); 7.00 (m, 2H); 6.85 (m, 1H); 3.94-3.73 (m, 4H); 3.66 (dd, J = 14,7,1H); 3.43 (dd, J = 14,6, 1H); 3.23-3.09 (m, 4H); 2.96-2.84 (m, 1H); 2.84 (s, 3H); 2.27-2.13 (m, 1H); 2.09-1.95 (m, 2H); 1.90-1.48 (m, 6H); 1.35 (s, 3H); 1.34 (s, 3H);

MS: 499 (M)<sup>+</sup>.

Example 19

In a manner analogous to that described in the first paragraph of Example 1, from 0.455 g of 4-[2(R)-[1-(R or S)-benzyloxycarbamoyl]-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-morpholine (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.194 g of 4-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]morpholine (diastereoisomer 1) in the form of a white solid.  
 nmr (MeOD): 3.80-3.51 (m, 9H); 3.42 (dd, J = 14,6,1H); 3.14-3.06 (dt, J = 4,11,1H); 3.04-2.86 (m, 1H); 2.85 (s, 3H); 2.23-2.11 (m, 1H); 2.06-1.95 (m, 2H); 1.91-1.73 (m, 2H); 1.71-1.46 (m, 4H); 1.35 (s, 3H); 1.34 (s, 3H);  
 MS:425 (M)<sup>+</sup>.

Example 20

In a manner analogous to that described in the first paragraph of Example 1, from 0.625 g of 1-[2(R)-[1-(R or S)-benzyloxycarbamoyl]-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-pyrrolidine (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.384 g of 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]morpholine (diastereoisomer 1) in the form of a white solid.  
 nmr (MeOD): 3.77-3.69 (m, 1H); 3.61 (dd, J = 14,6,1H); 3.53-3.44 (m, 2H); 3.39-3.31 (m, 2H); 2.93-2.85 (m, 2H); 2.84 (s, 3H); 2.26-2.13 (m, 1H); 2.07-1.71 (m, 8H); 1.69-1.46 (m, 4H); 1.36 (s, 3H); 1.33 (s, 3H);  
 MS:409 (M + H)<sup>+</sup>.

Example 21

In a manner analogous to that described in the first paragraph of Example 1, from 0.176 g of 8-[2(R)-[1-(R or S)-benzyloxycarbamoyl]-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-1,4-dioxo-8-azaspiro[4,5]decane (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.084 g of 8-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-1,4-dioxo-8-azaspiro[4,5]decane (diastereoisomer 1) in the form of a white solid.  
 nmr (MeOD): 4.02 (s, 4H); 3.81-3.60 (m, 5H); 3.99 (dd, J = 14,6,1H); 3.20-3.10 (m, 1H); 2.93-2.85 (m, 1H); 2.84 (s, 3H); 2.21-2.09 (m, 1H); 2.06-1.93 (m, 2H); 1.80-1.46 (m, 10H); 1.35 (s, 3H); 1.33 (s, 3H);  
 MS:481 (M + H)<sup>+</sup>.

Example 22

In a manner analogous to that described in the first paragraph of Example 1, from 0.443 g of 1-[2(R)-[1-(R or S)-benzyloxycarbamoyl]-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-4-methoxypiperidine (diastereoisomer 1) there was obtained 0.319 g of 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]-4-methoxypiperidine (diastereoisomer 1) in the form of a white solid.  
 nmr (MeOD): 3.96-3.80 (m, 2H); 3.69-3.59 (m, 1H); 3.54-3.23 (m, 7H); 3.18-3.09 (m, 1H); 2.93-2.80 (m, 4H); 2.21-2.09 (m, 1H); 2.07-1.41 (m, 12H); 1.41-1.38 (m, 6H);  
 MS:453 (M + H)<sup>+</sup>.

The starting material was prepared as follows

(i) A solution of 0.925 g of 1-[2(R)-[1(R or S)-(tert.butoxycarbonyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclopropylpropionyl]-4-hydroxypiperidine in 8 ml of dimethylformamide was treated with 1.08 g of methyl iodide and 1.79 g of silver oxide. The mixture was stirred at room temperature in the dark for 2 days. Additional portions of 0.54 g of methyl iodide and 0.895 g of silver oxide were then added and the mixture was stirred for a further 3 days. The solvent was evaporated and the residue was suspended in ethyl acetate and filtered. The ethyl acetate solution was concentrated and the residue was purified by flash chromatography on silica gel using ethyl acetate for the elution. There was obtained 0.61 g of 1-[2(R)-[1(R or S)-(tert.butoxycarbonyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-4-methoxypiperidine in the form of a colourless gum.

(ii) In a manner analogous to that described in Example 1 (viii)-(ix) from 0.61 g of 1-[2(R)-[1(R or S)-(tert.butoxycarbonyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-4-methoxypiperidine there was obtained 0.443 g of 1-[2(R)-[1(R or S)-(benzyloxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-4-methoxypiperidine (diastereoisomer 1)



in the form of a colourless gum.

#### Example 23

5 In a manner analogous to that described in the first paragraph of Example 1, from 0.94 g of 1-[2(R)-[1-(RS)-benzyloxycarbonyl]-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-octahydroazocine (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.663 g of 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]octahydroazocine (diastereoisomer 1) in the form of a white solid:  
 10 nmr (MeOD): 3.77 (dd, J = 14,10,1H); 3.66-3.43 (m, 4H); 3.33 (dd, J = 14,5,1H); 3.07 (dt, J = 10,4,1H); 2.91-2.81 (m, 4H); 2.29-2.16 (m, 1H); 2.10-1.95 (m, 2H); 1.90-1.46 (m, 16H); 1.34 (s, 6H); MS:451 (M + H)<sup>+</sup>.

#### Example 24

15 In a manner analogous to that described in the first paragraph of Example 1, from 0.37 g of 1-[2(R)-[1(R or S)-benzyloxycarbamoyl]-2-(5,5-dimethyl-2,4-dioxo-3-oxazolidinyl)ethyl]-3-cyclobutylpropionyl]piperidine (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (v)-(ix) using 3-(bromomethyl)-5,5-dimethyloxazolidine-2,4-dione in place of 1-(bromomethyl)-3,4,4-trimethyl-2,5-imidazolidinedione, there was obtained 0.131 g of 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(5,5-dimethyl-2,4-dioxo-3-oxazolidinyl)ethyl]propionyl]piperidine (diastereoisomer 1) in the form of a white solid:  
 20 nmr (MeOD): 3.72-3.53 (m, 5H); 3.39 (dd, J = 14,6,1H), 3.14 (dt, J = 10,4,1H); 2.95-2.86 (m, 1H); 2.23-2.11 (m, 1H); 2.08-1.94 (m, 2H); 1.90-1.44 (m, 18H); MS:410 (M + H)<sup>+</sup>.

#### Example 25

In a manner analogous to that described in the first paragraph of Example 1, from 0.42 g of 1-[2(R)-[1(R or S)-benzyloxycarbamoyl]-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-3-cyclobutylpropionyl]-  
 30 hexahydroazepine (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), there was obtained 0.197 g of 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]propionyl]hexahydroazepine (diastereoisomer 1) in the form of a white solid:  
 nmr (MeOD): 3.77-3.64 (m, 2H); 3.62-3.45 (m, 3H); 3.33 (dd, J = 14,5,1H); 3.07 (dt, J = 10,4,1H); 2.91-2.81 (m, 4H); 2.24-2.13 (m, 1H); 2.09-1.95 (m, 2H); 1.90-1.47 (m, 14H); 1.35 (s, 3H); 1.34 (s, 3H);  
 35 MS:437 (M + H)<sup>+</sup>.

#### Example 26

In a manner analogous to that described in the first paragraph of Example 1, from 0.37 g of 1-[2(R)-[1(R or S)-benzyloxycarbamoyl]-2-(hexahydro-1,3-dioxypyrazolo[1,2-a][1,2,4]triazol-2-yl)ethyl]-3-cyclobutyl-  
 40 propionyl]piperidine (diastereoisomer 1), prepared in a manner analogous to that described in Example 1 (i)-(ix), using 2-(bromomethyl)-hexahydro-1,3-dioxypyrazolo[1,2-a][1,2,4]triazole, there was obtained 0.118 g of 1-[3-cyclobutyl-2(R)-[2-(hexahydro-1,3-dioxypyrazolo[1,2-a][1,2,4]triazol-2-yl)-1(R or S)-(hydroxycarbamoyl)-ethyl]propionyl]piperidine in the form of a white solid.  
 45 nmr (MeOD): 3.68-3.56 (m, 8H); 3.52-3.39 (m, 2H); 3.17-3.09 (m, 1H); 2.97-2.90 (m, 1H); 2.35-2.27 (m, 2H); 2.21-2.11 (m, 1H); 2.07-1.95 (m, 2H); 1.88-1.44 (m, 12H); MS:422 (M + H)<sup>+</sup>.

#### Example 27

50 In a manner analogous to that described in the first paragraph of Example 1, from 0.222 g of 1-[2(R or S)-(benzyloxycarbamoyl)-2-phthalimidoethyl]-3-cyclobutylpropionyl]piperidine prepared in a manner analogous to that described in Example 1 (i)-(ix) using N-(bromomethyl)-phthalimide, there was obtained 0.013 g of 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-phthalimidoethyl]propionyl]piperidine (diastereoisomer 1) in the form of a white solid.  
 55 nmr (MeOD): 7.87-7.75 (m, 4H); 3.83 (dd, J = 14.8, 1H); 3.66-3.58 (m, 3H); 3.53-3.45 (m, 1H); 3.35-3.25 (m, 1H); 3.20-3.12 (m, 1H); 3.04-2.97 (m, 1H); 2.23-2.11 (m, 1H); 2.08-1.95 (m, 2H); 1.89-1.41 (m, 12H); MS:428 (M + H)<sup>+</sup>.

# EP 0 684 240 A1

The following Examples illustrate pharmaceutical preparations containing the hydroxamic acid derivatives provided by the present invention:

## Example A

Tablets containing the following ingredients may be produced in a conventional manner:

Ingredient	Per tablet
Hydroxamic acid derivative	10.0 mg
Lactose	125.0 mg
Corn starch	75.0 mg
Talc	4.0 mg
Magnesium stearate	1.0 mg
Total weight	215.0 mg


## Example B

Capsules containing the following ingredients may be produced in a conventional manner:

Ingredient	Per capsule
Hydroxamic acid derivative	10.0 mg
Lactose	165.0 mg
Corn starch	20.0 mg
Talc	5.0 mg
Capsule fill weight	200.0 mg

## Claims

### 1. Compounds of the general formula



wherein

R<sup>1</sup> represents cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl;

R<sup>2</sup> represents a saturated 5- to 8-membered monocyclic or bridged N-heterocyclic ring which is attached via the N atom and which, when it is monocyclic, optionally contains NR<sup>4</sup>, O, S, SO or SO<sub>2</sub> as a ring member and/or is optionally substituted on one or more C atoms by hydroxy, lower alkyl, lower alkoxy, oxo, ketalized oxo, amino, mono(lower alkyl)amino, di-(lower alkyl) amino, carboxy, lower alkoxy-carbonyl, hydroxymethyl, lower alkoxy-methyl, carbamoyl, mono(lower alkyl)-carbamoyl, di(lower alkyl)carbamoyl or hydroxyimino;

R<sup>3</sup> represents a 5- or 6-membered N-heterocyclic ring which (a) is attached via the N atom, (b)

optionally contains N, O and/or S, SO or SO<sub>2</sub> as a additional ring member, (c) is substituted by oxo on one or both C atoms adjacent to the linking N atom and (d) is optionally benz-fused or optionally substituted on one or more other C atoms by lower alkyl or oxo and/or on any additional N atom(s) by lower alkyl or aryl;

5 R<sup>4</sup> represents hydrogen, lower alkyl, aryl, aralkyl or a protecting group;

m stands for 1 or 2; and

n stands for 1-4;

and pharmaceutically acceptable salts thereof.

10 2. Compounds according to claim 1, wherein

R<sup>1</sup> represents cyclopropyl, cyclobutyl or cyclopentyl;

15 R<sup>2</sup> represents a 5-, 6- or 7-membered monocyclic or bridged N-heterocyclic ring which is attached via the N atom and which, when it is monocyclic, optionally contains NR<sup>4</sup>, O, S, SO or SO<sub>2</sub> as a ring member and/or is optionally substituted on one or more C atoms by hydroxy, lower alkyl, lower alkoxy, oxo, ketalized oxo, amino, mono(lower alkyl)amino, di-(lower alkyl) amino, carboxy, lower alkoxy carbonyl, hydroxymethyl, lower alkoxy methyl, carbamoyl, mono(lower alkyl)-carbamoyl, di(lower alkyl)carbamoyl or hydroxyimino;

20 R<sup>3</sup> represents a 5- or 6-membered N-heterocyclic ring which (a) is attached via the N atom, (b) optionally contains N, O and/or S as an additional ring member in a position or positions other than adjacent to the linking N atom, (c) is substituted by oxo on one or both C atoms adjacent to the linking N atom and (d) is optionally benz-fused or optionally substituted on one or more other C atoms by lower alkyl or oxo and/or on any additional N atom(s) by lower alkyl or aryl; and

25 R<sup>4</sup> represents hydrogen, lower alkyl or a protecting group.


3. Compounds according to claim 1 or claim 2, wherein R<sup>2</sup> represents 1-pyrrolidinyl, piperidino, 4-aryl-1-piperazinyl, morpholino, tetrahydro-1,4-thiazin-4-yl, tetrahydro-1,4-thiazin-4-yl 1,1-dioxide, thiazolidin-3-yl, hexahydroazepino or octahydroazocino optionally substituted on one or more C atoms by hydroxy, lower alkyl, lower alkoxy, ketalized oxo or mono(lower alkyl)-carbamoyl; or 3-azabicyclo[3.2.2]nonane.

30

4. Compounds according to claim 3, wherein R<sup>2</sup> represents piperidino or hydroxypiperidino.

5. Compounds according to claim 4, wherein hydroxypiperidino is 4-hydroxypiperidino.

35 6. Compounds according to any one of claims 1 to 5, wherein R<sup>3</sup> represents a group of the formula



50 wherein

R<sup>5</sup> and R<sup>6</sup> each represent hydrogen or together represent an additional bond or the remainder of a fused benzene ring; R<sup>7</sup> represents hydrogen, lower alkyl or aryl;

X represents -CO-, -CH<sub>2</sub>-, -CH(lower alkyl)-, -C(lower alkyl)<sub>2</sub>-, -NH-, -N(lower alkyl)- or -O-; or, when R<sup>7</sup> represents lower alkyl and X represents -N(lower alkyl)-, the lower alkyl groups can be joined to form a 5-, 6- or 7-membered ring;

55

7. Compounds according to claim 6, wherein R<sup>3</sup> represents a group of formula c), R<sup>7</sup> represents lower alkyl and X represents -C(lower alkyl)<sub>2</sub>-.

8. Compounds according to claim 7, wherein R<sup>3</sup> represents 3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl.
9. Compounds according to any one of claims 1 to 8, wherein m and n stand for 1.
10. 1-[3-Cyclopropyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]piperidine.
11. 1-[3-Cyclopropyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]-4-piperidinol.
12. 1-[3-Cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]piperidine.
13. 1-[3-Cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-4-piperidinol.
14. 1-[3-Cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]-4-piperidinol.
15. 1-[3-Cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)ethyl]-propionyl]piperidine.
16. A compound according to claim 2, selected from:  
 3-[3-Cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]-3-azabicyclo[3.2.2]nonane,  
 3-[3-cyclopropyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]-3-azabicyclo[3.2.2]nonane and  
 3-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]-3-azabicyclo[3.2.2]nonane.
17. A compound according to claim 1, selected from:  
 1-[3-cyclohexyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]piperidine,  
 4-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]tetrahydro-1,4-thiazine,  
 4-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]tetrahydro-1,4-thiazine S,S-dioxide,  
 4-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]tetrahydro-1,4-thiazine,  
 4-[3-cyclohexyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]tetrahydro-1,4-thiazine,  
 3-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]-5,5-dimethyl-N-propyl-4(R)-thiazolidinecarboxamide,  
 4-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]morpholine,  
 3-[3-cyclopentyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]-N,5,5-trimethyl-4(R)-thiazolidinecarboxamide,  
 4-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]-4-phenylpiperazine,  
 4-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]morpholine,  
 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]pyrrolidine,  
 8-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]-1,4-dioxo-8-azaspiro[4.5]decane,  
 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]-4-methoxypiperidine,  
 1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-

ethyl]propionyl]octahydroazocine,


1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(5,5-dimethyl-2,4-dioxo-3-oxazolidinyl)ethyl]-propionyl]piperidine,

1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-(3,4,4-trimethyl-2,5-dioxo-1-imidazolidinyl)-ethyl]propionyl]hexahydroazepine,

1-[3-cyclobutyl-2(R)-[2-(hexahydro-1,3-dioxypyrazolo[1,2-a][1,2,4]triazol-2-yl)-1(R or S)-(hydroxycarbamoyl)ethyl]propionyl]piperidine and


1-[3-cyclobutyl-2(R)-[1(R or S)-(hydroxycarbamoyl)-2-phthalimidoethyl]propionyl]piperidine.

18. Compounds of the general formula



wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, m and n have the significance given in claim 1.

19. Compounds of the general formula




wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, m and n have the significance given in claim 1.


20. Compounds according to any one of claims 1-17 for use as therapeutically active substances, especially in the control or prevention of degenerative joint diseases or in the treatment of invasive tumours, atherosclerosis or multiple sclerosis.

21. A process for the manufacture of the compounds claimed in any one of claims 1-17, which process comprises


(a) reacting an acid of the general formula



15 wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $m$  and  $n$  have the significance given in claim 1, with a compound of the general formula



wherein  $Z$  represents hydrogen, tri(lower alkyl)silyl or diphenyl(lower alkyl)silyl, and, where required, cleaving off any diphenyl(lower alkyl)silyl group present in the reaction product, or  
(b) catalytically hydrogenating a compound of the general formula



40 wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $m$  and  $n$  have the significance given in claim 1 and  $Bz$  represents benzyl, and,  
if desired, converting a compound of formula I obtained into a pharmaceutically acceptable salt.

45 22. A process for the manufacture of medicaments, especially for use in the control or prevention of degenerative joint diseases or in the treatment of invasive tumours, atherosclerosis or multiple sclerosis, which process comprises bringing a compound according to any one of claims 1-17 with a therapeutically inert carrier material and bringing the mixture into a galenical administration form.

50 23. A medicament containing a compound according to any one of claims 1-17 and a therapeutically inert carrier material.

24. A medicament for the control or prevention of degenerative joint diseases or for the treatment of invasive tumours, atherosclerosis or multiple sclerosis, containing a compound according to any one of claims 1-17 and a therapeutically inert carrier material.

55 25. The use of a compound according to any one of claims 1-17 in the control or prevention of illnesses, especially for the control or prevention of degenerative joint diseases or for the treatment of invasive tumours, atherosclerosis or multiple sclerosis.

**EP 0 684 240 A1**

**26.** The use of a compound according to any one of claims 1-17 for the manufacture of a medicament for the control or prevention of degenerative joint diseases or for the treatment of invasive tumours, atherosclerosis or multiple sclerosis.

5   **27.** Compounds according to any one of claims 1-17, whenever prepared according to a process according to claim 21.

10

15

20

25

30

35

40

45

50

55



European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 95 10 5495  
shall be considered, for the purposes of subsequent  
proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL. 6)
Y	EP-A-0 574 758 (F. HOFFMANN-LA ROCHE A.G.) * the whole document * ---	1,23	C07D401/06 C07D403/06 C07D413/06 C07D417/06 C07D487/04 A61K31/445 C07C259/06
Y	EP-A-0 575 844 (F. HOFFMANN-LA ROCHE A.G.) * claim 1 * ---	1,23	
Y	EP-A-0 489 577 (CELLTECH LTD.) * example 1 * ---	1,23	
A,P	WO-A-94 22309 (GLYCOMED INC.) * examples * ---	1,23	
A	WO-A-91 02716 (BRITISH BIO-TECHNOLOGY LTD.) * examples 13-17 * ---	1,23	
A	WO-A-90 05716 (BRITISH BIO-TECHNOLOGY LTD.) * ex. 11,18,19,23-26,28,31-36,38 * ---	1,23	
A,P	WO-A-95 04715 (KANEBO LTD.) * abstract * --- -/--	1,23	TECHNICAL FIELDS SEARCHED (Int. CL. 6)  C07D A61K C07C
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims</p> <p>Claims searched completely: Claims searched incompletely: Claims not searched: Reason for the limitation of the search:</p> <p>Remark: Although claim 25 is directed to a method of treatment of (diagnostic method practised on) the human/animal body (Article 52(4) EPC) the search has been carried out and based on the alleged effects of the compound/composition.</p>			
Place of search BERLIN		Date of completion of the search 18 September 1995	Examiner Frelon, D
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document	





European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 95 10 5495

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A,P	EP-A-0 613 883 (YAMANOUCI PHARMACEUTICAL CO., LTD.) * the whole document * ---	1,23	
A	EP-A-0 405 788 (SHIONOGI SEIYAKU KABUSHIKI KAISHA) * example 5; tables 1-2 * -----	1,23	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)